

**AQUAPONIC BIOFILTER SYSTEM USING *HELICONIA* SP. FOR AMMONIA  
REMOVAL IN LAKE**

**MUHAMMAD SAFWAN BIN MISWAN**

A thesis submitted in partial  
fulfillment of the requirement for the award of the  
Degree of Master of Civil Engineering

Faculty of Civil and Environmental Engineering

Universiti Tun Hussein Onn Malaysia

MARCH 2020

بِسْمِ اللَّهِ الرَّحْمَنِ الرَّحِيمِ  
السلام عليكم

SPECIALLY DEDICATED TO MY ADORED  
PARENTS, MY WIFE, FAMILY AND FRIEND FOR  
ALWAYS BEING BESIDE ME. THANKS FOR ALL  
THE PATIENCE, SUPPORT AND LOVE.  
SEMOGA ALLAH RAHMATI KAMU SEMUA



## ACKNOWLEDGEMENT

*With the name of Allah the Most Merciful and His messenger Prophet*

*Muhammad p.b.u.h*

Alhamdulillah, with the blessing of Allah, the study have been successfully completed. I would like to take this opportunity to extend my appreciation and thankfulness to my supervisor, Prof. Emeritus IR. DR. Amir Hashim Bin Mohd Kassim and my co-supervisors, Assoc. Prof. TS. Dr. Radin Maya Saphira Binti Radin Mohamed and TS. DR. Adel Ali Saeed Abduh Algheethi for their kind guidance and support toward the accomplishment of my master degree journey.

Many thanks also to all assistant engineer in FKAAS for their friendly assistance and help. Their views and tips are very useful indeed to complete my study. Thanks for all my friends for their support, help and for being there when i need them throughout the year for solving over to finish my study. v

Not forgetting also, i would like to express my gratitude and love to my loving parents, my such caring wife and my back-bone family members for their guidance and supports in my study. May Allah reward all yours goodness. Thank you.

*May Allah showering His blessing upon us al*

## ABSTRACT

Lakes are considered as one of the natural water resources in Malaysia. The quick pace of development around many lakes catchment areas has a significant impact on water quality. Consequently, biofilter aquaponic system for ammoniacal nitrogen removal in lake near non-point sources such as oil palm plantation, factory and educational building were studied. Aquaponic is a system that mutually integrates aquaculture and plant cultivation. Aquaponic biofilter system also applies phytoremediation mechanism to reduce nutrient. Therefore, phytoremediation of lake water in *Tasik Teknologi* in campus area of Universiti Tun Hussein Onn Malaysia (UTHM) was performed in 7 days by using *Heliconia* sp. as phytoremediate plant. This study was conducted in a greenhouse experiment to control the environments conditions (with different temperature and photoperiod), which consists of five hydroponic tanks, and one fish culture tank with the 40 watt of water pump was run circulation of water. A physicochemical characteristic of *Tasik Teknologi* was compared to Interim National Water Quality Standard Malaysia (NWQSM) after phytoremediation. *Heliconia* sp. after phytoremediation was undergoes analysis for plant nutrients by using Ion Chromatography (IC) in three major parts; leaves, roots and stem. The result shows that *Tasik Teknologi* has mean values of AN (3.95 mg/L), NO<sub>2</sub> (0.12 mg/L), NO<sub>3</sub> (1.36 mg/L), BOD (105.70 mg/L), COD (10.30 mg/L), TSS (19.70 mg/L), Turbidity (12.30 mg/L), temperature (28.80°C), DO (9.40 mg/L) and pH (6.51), respectively. Phytoremediation has reduced AN and NO<sub>3</sub> from the lake water sample by *Heliconia* sp. with 66.00% and 92.03% removal efficiency, respectively. Besides that, the concentration of DO during phytoremediation was increased 47.82 % from 2.4 mg/L to 4.6 mg/L. Furthermore, the uptake of AN in three major parts of *Heliconia* sp. leaves, stem and root were found to be 4.40 %, 1.12% and 4.20%, respectively. The *Heliconia* sp. has the ability in reducing AN and NO<sub>3</sub> in lake water with efficiently and practically.

## ABSTRAK

Tasik merupakan sumber semulajadi dan air yang penting di Malaysia. Kadar yang cepat untuk pembangunan di seluruh kawasan tadahan tasik mempunyai kesan yang ketara terhadap kualiti air. Oleh itu, sistem penapis-bio akuaponik untuk penyingkiran ammoniakal nitrogen di tasik berhampiran bukan titik punca seperti ladang kelapa sawit, kilang dan bangunan pendidikan telah dikaji. Akuaponik adalah satu sistem yang saling mengintegrasikan akuakultur dan penanaman tumbuhan. Akuaponik penapis-bio sistem juga menggunakan mekanisme fitopemuliharaan untuk mengurangkan nutrien. Oleh itu, fitopemuliharaan air tasik di Tasik Teknologi di kawasan Kampus Universiti Tun Hussein Onn Malaysia (UTHM) telah dijalankan selama 7 hari dengan menggunakan pokok *Heliconia* sp. sebagai pokok fitopemuliharaan. Kajian ini dijalankan dalam satu eksperimen rumah hijau untuk mengawal keadaan persekitaran (dengan pelbagai bacaan suhu dan foto masa), yang terdiri daripada lima tangki hidroponik, dan satu tangki ikan dengan 40 watt pam air untuk menjalankan peredaran air. Ciri fizikokimia Tasik Teknologi telah dibandingkan dengan Piawaian Kualiti Air Negara Malaysia (NWQSM) selepas proses fitopemuliharaan. *Heliconia* sp. selepas proses fitopemuliharaan terus di analisis nutrien tumbuhan dengan menggunakan ion kromatografi (IC) dalam tiga bahagian utama; daun, akar dan batang. Hasilnya menunjukkan bahawa Tasik Teknologi mempunyai nilai AN (3.95 mg/L), NO<sub>2</sub> (0.12 mg/L), NO<sub>3</sub> (1.36 mg/L), BOD (105.70 mg/L), COD (10.30 mg/L), TSS (19.70 mg/L), kekeruhan (12.30 mg/L), suhu (28.80 °C), DO (9.40 mg/L) dan pH (6.51). Fitopemuliharaan telah mengurangkan AN dan NO<sub>3</sub> dari sampel air Tasik oleh *Heliconia* sp. sebanyak 66.00% dan 92.03% dengan kecekapan penyingkiran. Di samping itu, larutan DO semasa fitopemuliharaan meningkat sebanyak 47.82% daripada 2.4 mg/L kepada 4.6 mg/L. Tambahan pula, penyerapan AN dalam tiga bahagian utama *Heliconia* sp. iaitu daun, batang dan akar adalah 4.40%, 1.12% dan 4.20%. *Heliconia* sp. mempunyai keupayaan untuk mengurangkan AN dan NO<sub>3</sub> dalam air tasik dengan cekap dan praktikal.

## TABLE OF CONTENTS

<b>TITLE</b>	<b>i</b>
<b>DECLARATION</b>	<b>ii</b>
<b>DEDICATION</b>	<b>iii</b>
<b>ACKNOWLEDGEMENT</b>	<b>iv</b>
<b>ABSTRACT</b>	<b>v</b>
<b>ABSTRAK</b>	<b>vi</b>
<b>TABLE OF CONTENTS</b>	<b>vii</b>
<b>TABLE OF FIGURES</b>	<b>viii</b>
<b>TABLE OF TABLES</b>	<b>xiii</b>
<b>LIST OF SYMBOLS AND ABBREVIATION</b>	<b>xv</b>
<b>CHAPTER 1 INTRODUCTION</b>	<b>1</b>
1.1 Background of Study	1
1.2 Problem Statement	4
1.3 Objectives of Study	6
1.4 Scope of Study	6
1.5 Hypotheses of the Study	7
1.6 Significance of Study	8
1.7 Thesis structure	8
<b>CHAPTER 2 LITERATURE REVIEW</b>	<b>10</b>
2.1 Introduction	10
2.2 Lake Water in Malaysia and Water Quality Parameter Standard	10
2.3 Conventional Nitrogen, Ammoniacal nitrogen and Nitrate Removal Technologies	17

2.4	Aquaponic Biofilter System (ABS) in Malaysia	21
2.5	Phytoremediation and its Efficiency	28
2.6	Selection of Tilapia ( <i>Oreochromis niloticus</i> ) for Aquaponic Biofilter System	31
2.7	Plant Applications for Ammoniacal Nitrogen Removal in Aquaponic Biofilter System	34
2.8	Temperature Effect on Plant Growth in Aquaponic Biofilter System	35
2.9	Photoperiod Effect on Plant Growth in Aquaponic Biofilter System	36
2.10	Nutrient Removal and the mechanism of nitrification	36
2.11	Previous Studies on Biofilter and its Ability to Remove Pollutants	39
2.12	Key Findings of Literature Review	42
<b>CHAPTER 3</b>	<b>METHODOLOGY</b>	<b>43</b>
3.1	Introduction	43
3.2	Outlines of Methodology	43
3.3	Overview Methodology Flow Chart	44
3.4	Study Area	45
3.5	Lake Water Sampling	46
3.6	Sampling Preservation and Storage	47
3.7	Characterization of <i>Tasik Teknologi</i> in UTHM	48
3.8	Determination of Ammonia Nitrogen (AN)	49
3.9	Hydroponic Tanks and Cultivation of Plant	49
3.10	Fish Culture Tank and Acclimation	50

3.11	Design and Cycle of Aquaponic Biofilter System	51
3.12	Experimental Procedure of Aquaponic Biofilter System	53
3.13	Water Quality	55
3.14	Determination Aquaponic Biofilter System performances	55
3.15	Aquaponic Biofilter System And Its Efficiency	56
3.16	Data Analysis	57

## **CHAPTER 4      RESULT AND DISCUSSION      58**

4.1	Introduction	58
4.2	<i>Tasik Teknologi</i> Characteristics	58
4.3	Characteristics of <i>Heliconia</i> sp. in Aquaponic Biofilter System on Floating Bed	64
4.4	Acclimatization Phase of the Aquaponics Biofilter System	65
4.5	Efficiency of Aquaponics Biofilter System for Ammonia, Nitrite and Nitrate Removal	66
4.6	Influence of the Hydraulic Retention Time (HRT) and Hydraulic Loading Rate (HLR)	71
4.7	Correlation between pH and DO with ammonia in aquaponics biofilter system after 7 days of phytoremediation process	73
4.8	The correlation between ammoniacal nitrogen and nitrate with root of <i>Heliconia</i> sp. growth as fertilizer for the plant	76
4.9	The concentration ammonia , nitrate and nitrite in root, stem and leaves of <i>Heliconia</i> sp .using Ion Chromatography (IC)	78



4.9.1	The concentration of ammonia nitrite and nitrate before and after treatment after 7 days in leaves of <i>Heliconia</i> sp.	79
4.9.2	The concentration ammonia and nitrate before and after treatment after 7 days of phytoremediation in stem of <i>Heliconia</i> sp.	81
4.9.3	The concentration before and after treatment after 7 days in root of <i>Heliconia</i> sp.	83
4.10	The key finding of research outcomes	85
<b>CHAPTER 5</b>	<b>CONCLUSION AND RECOMMENDATION</b>	<b>89</b>
5.1	Introduction	87
5.2	Conclusions	87
5.3	Recommendations	88
<b>REFERENCES</b>		<b>90</b>
<b>APPENDIX</b>		<b>110</b>



## LIST OF TABLES

2.1	Lakes and Reservoirs Based on States in Malaysia	11
2.2	Interim National Water Quality Standards for Malaysia	13
2.3	Water Quality Classification Based on Water Quality Index (WQI)	14
2.4	Parameters of Water Quality Standard Based on Environmental Protection Agency	14
2.5	Characteristics of Lake Water by Previous Studies in Malaysia	15
2.6	The Advantages and Disadvantages of Nitrogen Removal Technologies	19
2.7	Type of Aquaponics and Their Efficiency Reported in the Literature in Malaysia, Taiwan and United States	28
2.8	Different Phytoremediation Mechanisms used to Remediate Varying Pollutants.	30
2.9	The Retention Time for Pollutant Removal by Phytoremediation	31
2.10	Culture Condition of Tilapia ( <i>Oreochromis niloticus</i> ) in Previous Studies	33
2.11	The Previous Studies on <i>Heliconia</i> sp. in Remove Pollutants	35

2.12	The Previous Studies of Constructed Wetland in Nitrogen Removal	41
3.1	Sample Preservation and Storage	48
3.2	The Physicochemical Characteristics Measurement on <i>Tasik Teknologi</i> Water UTHM	48
4.1	The geometrical and environmental conditions of <i>Tasik Teknologi</i> UTHM	59
4.2	Characteristics Of <i>Tasik Teknologi</i> Water in UTHM Campus Area With Minimum, Maximum and Mean Value $\pm$ Standard Deviation. Sampling Was Conducted On March 2018 And September 2018. (n=12, Mean $\pm$ SD)	60
4.3	Classification of Ten Water Quality Parameters in <i>Tasik Teknologi</i> UTHM According to NWQS Malaysia Standard	64
4.4	Initial Physical Characteristics of <i>Heliconia</i> sp. in the Aquaponics Biofilter System	65
4.5	Control Condition Aquaponic Biofilter System during 7 days of Phytoremediation	66
4.6	Average and standard deviation of ammoniacal nitrogen (AN) during 7 days of treatment	67
4.7	Average and standard deviation of water quality parameter in term of Nitrate (NO <sub>3</sub> ) after 7 days of treatment	69
4.8	The percentage removal of Ammoniacal Nitrogen and Nitrate in each parameter compare to the previous researcher	70
4.9	The design feature of Aquaponic Biofilter System with percentage of removal for ammonia and nitrate	72
4.10	The concentration of before and after ammonia, nitrite and nitrate in leaves of <i>Heliconia</i> sp.	79

4.11	The concentration of ammonia and nitrate in stem of <i>Heliconia</i> sp. for each parameter	81
4.12	The concentration of ammonia and nitrate in root of <i>Heliconia</i> sp. after 7 days phytoremediation	84



## LIST OF FIGURES

2.1	The Production of Aquaculture in Malaysia from Years 2011 to 2017	23
2.2	Publication on Aquaculture and Aquaponic in Malaysia	24
2.3	The Media Filled System Schematic Diagram	25
2.4	The Nutrient Film Technique Schematic Diagram	26
2.5	The Schematic Diagram of Floating Raft System for this study	27
2.6	Nitrification Process and Important Bacteria	38
3.1	Methodology Flow Chart	44
3.2	Study Area at Universiti Tun Hussein Onn Malaysia (UTHM)	45
3.3	Location of Lake near Oil Palm Plantation in UTHM	46
3.4	Schematic Diagram of Sampling Point and Depth of Sampling	47
3.5	The Algae and Plant Growth Occur on Surface of Water	47
3.6	<i>Heliconia psittacorum</i> sp.	50
3.7	Design of Hydroponic Tank and Cultivation of Plant	50
3.8	Schematic Design of Aquaponic Biofilter System	52
4.1	The height of maximum and minimum of <i>Heliconia</i> sp.	65
4.2	The trend of ammonia removal after 7 days in every parameters	68
4.3	The percentage of Nitrate removal during 7 days in every parameters	69

4.4	The HRT and HLR in the present study compare to previous study in affect percentage removal of ammonia and nitrate	73
4.5	Water quality parameter in Aquaponic Biofilter System before and after phytoremediation in 7 days	74
4.6	The correlation between demand oxygen (DO) with ammonia along 7 days phytoremediation	74
4.7	The correlation between pH and ammonia along 7 days phytoremediation	75
4.8	The regeneration of new root after 7 days of phytoremediation in Aquaponic Biofilter Sysrem before and after treatment	77
4.9	Pattern showing the increases in size of surface area of roots against age, and correlation between the size surface area of roots <i>Heliconia</i> sp., ammonia and nitrate	78
4.10	The graph pattern of ammonia, nitrite and nitrate concentration before and after treatment after 7 days in leaves of <i>Heliconia</i> sp.	80
4.11	The graph of ammonia and nitrate concentration in Aquaponic Biofilter System	82
4.12	The concentration of ammonia and nitrate in root after 7 days phytoremediation	84

## LIST OF SYMBOLS AND ABBREVIATIONS

°C	Degree Celcius
ANOVA	Analysis of Variance
ABS	Aquaponic Biofilter System
AOB	Ammonia oxidizing bacteria
BOD	Biochemical Oxygen Demand
cm	Centimeter
COD	Chemical Oxygen Demand
DO	Dissolved Oxygen
DOE	Department of Environment
EPA	Environmental Protection Agency
FAO	Food Agriculture Organization
HRT	Hydraulic Retention Time
HLR	Hydraulic Loading Rate
INWQS	Interim National Water Quality Standards
mg/L	Miligram per liter
MPOB	Malaysia Palm Oil Board
N	Nitrogen
AN	Ammoniacal Nitrogen
NH <sub>3</sub>	Ammonia
NO <sub>3</sub> <sup>-</sup>	Nitrate
NO <sub>2</sub> <sup>-</sup>	Nitrite
NOB	Nitrite oxidizing bacteria
P	Phosphorus
POME	palm oil mill effluent
TN	Total nitrogen
TSS	Total suspended solid

UTHM      Universiti Tun Hussein Onn Malaysia

WQI      Water Quality Index





## CHAPTER 1

### INTRODUCTION

#### 1.1 Background of study

Malaysia is one of the countries that are bestowed with biodiversity from all sources including rivers, lakes and sea. Rivers and lakes are natural heritages that should be preserved at all cost. The history of human civilization has proven the importance of rivers and lakes to humans but development activities have affected the quality status of these waterbodies. Pollution causes the waterbodies to lose its benefits to human and other living things. The pollution comes from point source (factories and sewage treatment plant) and non-point source (oil, fertilizers and lawn chemicals). This study focuses on palm oil plantation as non-point source and was conducted to evaluate the water quality changes in the lake of *Tasik Teknologi* at Universiti Tun Hussein Onn Malaysia (UTHM), Malaysia. Hence, the background study of palm oil plantation widely well known in Malaysia since a few years ago.

Palm oil plantations make up approximately 77% of agricultural land in Malaysia (Nadzir *et al.*, 2019). Palm oil plantation is broadly established in tropical countries such as Malaysia, Indonesia, and Thailand for its commercial oil. The Malaysia Palm Oil Board (MPOB) stated that the palm oil plantation area in Malaysia reached 5.85 million

hectares in 2018, an increase of 0.7% from 5.81 million hectares in 2017 (MPOB, 2018). Sarawak is the largest state with palm oil planted area in Malaysia. It has 1.57 million hectares or 26.9% of the total palm oil planted area, followed by Sabah with 1.55 million hectares or 26.5%. In comparison, Peninsular Malaysia (with 11 states) is accounted for 2.73 million hectares or 46.6% of the total planted area. Johor is the second largest state with the palm oil plantation area of 747,562 hectares and an estimated 12.8% in peninsular Malaysia (MPOB, 2018). Therefore, the increasing in total plantation area affects the utilization of fertilizer. Fertilizer was used to increase the production of palm oil in Malaysia.

The plantation may pose significant environmental pollution from palm oil mill effluent (POME), fertilizers, pesticides, insecticides and herbicides (Abdullah & Sulaiman, 2013). The excessive utilization of fertilizers in palm oil plantation that runoff to the nearby water body can cause eutrophication due to the high loads of nitrogen (N) and phosphorus (P) (Al-Badaii, Shuhaimi & Gasim, 2013). The authors reported the concentration of N and P in river water contaminated from agricultural activities at 1.91 mg/L and 1.9 mg/L, respectively. POME has high nutrient concentrations and pollutants, so releasing it into a water body will lead to algal blooms and subsequent loss of biologically active oxygen in the water (Isabelle, 2017). The lack of oxygen will eventually cause a loss to all life in the aquatic ecosystems. In Madaki & Seng (2013), the N and P concentrations in lake water influenced by POME wastes were reported 750 mg/L and 180 mg/L, respectively. Both researchers found that the level of N and P were exceeded the level regulated by Interim National Water Quality Standard for Malaysia (NWQS) which are <0.1 mg/L and 0.2 mg/L, respectively. Nadzi *et al.* (2019) stated that the application of toxic agrochemicals such as phosphate fertilizer can contaminate the nearby surface water with some heavy metals, organic and inorganic matters. Thus, it will negatively impact aquatic species as well as drinking water sources. Aquaponic Biofilter System (ABS) is one of the potential treatment systems that can be applied to treat aquaculture wastewater. ABS use the combination of aquaculture and hydroponic systems, where (*Heliconia* plants) are planted in a recirculating system that utilizes less water than the traditional farming. Nutrients contained in fish tanks are recycled into plant biomass in the presence of nitrifying bacteria that will convert the excreted

ammoniacal nitrogen (AN) to nitrite and then to nitrate. The water in recirculating system will be clear from ammonia and others substances by using application of plants that potentially as green technology compare others conventional method. There are few researcher that interested in this system and the results from their studies were satisfactory. Lam *et al.* (2015) claimed that ABS was used to treat aquaculture pond. Endut *et al.* (2009) also used ABS to treat nutrient-rich wastewater by using water spinach. Lam *et al.* (2015) and Endut *et al.* (2009) revealed that 70% and 86 % of ammoniacal nitrogen (AN) had been removed, respectively, confirming the ability of ABS to reduce nutrient from wastewater. Estim *et al.* (2018) reported a removal of 59% of PO<sub>4</sub>-N and 50% of AN from aquaculture wastewater by an indoor aquaponic system. The satisfactory results of Estim *et al.* (2018) study could be due to the lack of photoperiod for photosynthesis process. However, previous studies have not investigated the detail on the optimisation study for the efficiency of ABS. Factors such as photoperiod and temperature of ABS in laboratory scale are found to be important which can determine the survival of the plant over a certain time. These two factors are identified to influence the growth of plant, the time to flower, and the performances on efficiency removal of nutrients. Ha *et al.* (2013) stated that photoperiod and temperature have been reported to affect floral initiation and flower development of flowering plant families.

There are many advantages of aquaponics such as reducing the need for formulated fertilizers, eliminating the possibility of agricultural run-off, and cleansing of water through biofilter treatments (Turcios & Papenbrock, 2014). Turkmen and Guner (2010) stated that there is no mixing fertilizer involved in aquaponic, making it a great way for beginners to cultivate plants. Nutrients excess in the lake can be treated with filtration method using plants. Plants act as biofilter to pollutant in water by the uptake of N and P from water for growth, thus improve the water quality. The nutrients released from fish excreta and microbial breakdown of organic wastes are used by plants in aquaponic systems (Yavuzcan *et al.*, 2017). In the system, the plant component serves as a biofilter; therefore, a separate biofilter is not needed unlike aquaculture systems. In addition, the biofilter can generates income through the sale of the economic plant products (Saha *et al.*, 2016). Hence, the aquaponic systems develop an economically advantageous symbiotic system, where aquatic organisms and the plant component benefit each other

and while the grower receives two valuable commercial products. In contrast, the crop plant is the only marketable product in aquaponics and it is devoid of the commercial aquatic species and associated nutrient supply.

Many plant species such as *Heliconia sp.* ornamental plants (*Heliconia psittacorum*) have been investigated for removal of nitrogen (ammoniacal nitrogen) in phytoremediation. Konnerup *et al.* (2009) claimed that *Heliconia sp.* removed 13% of total nitrogen from domestic wastewater containing 27 mg/L nitrogen. Mendoza *et al.* (2015) reported that *Heliconia sp.* removed 39% of total nitrogen from domestic wastewater. These two findings showed the low efficiency of *Heliconia sp.* to remove TN in domestic wastewater. The low ability of *Heliconia sp.* as biofilter in this system implies that sub-surface flow systems have limited use if nutrient removal is required. Plant uptake is a significant removal process for nutrients, and only in lightly loaded system can this be sufficient to secure low effluent concentrations. Conversely, Parra (2016) found that, *Heliconia sp.* successfully removed 72% of total nitrogen in landfill leachate due to the underground tissues (root) showed a greater accumulation capacity. However, there are few attempts of using the *Heliconia sp.* in removing of nutrients such as ammoniacal nitrogen, nitrate and nitrite from the lake water. Therefore, this study was conducted to investigate the efficiency of using *Heliconia sp.* in removal of ammoniacal nitrogen from lake water near palm oil plantation.

## 1.2 Problem statement

The widely used chemical fertilizers in palm oil plantations has been reported to release the excessive nutrients (N and P) into the nearby water body causing the concentration level to exceed the standard regulated level (0.1 mg/L and 0.2 mg/L). This exceeding level is normally associated with the eutrophication phenomenon. The phenomenon occurs due to the excess of nutrients, which are nitrogen and phosphorus that will stimulate the algae growth. Over 20 years ago, the estimated fertilizer used for palm oil industries (estate only) in Malaysia is about 280,133 and 315,148 tones for nitrogen and phosphorus, respectively (FAO, 2004). Nowadays, there are various.

types of conventional nitrogen removal technologies, which can be divided into physiochemical and biological means. Each method relies on different operating principles and conditions. However, these technologies are costly in terms of operation, maintenance and production because they require chemical reagents or complementary facilities (Mishra & Maiti, 2017).

ABS technology has the ability to remove nutrients from lake water. Previous researchers such as Marques *et al.* (2017) and Boxman *et al.* (2018) suggested using an environmental friendly technology such as aquaponic system which is used as an efficient nutrients remover. Aquaponic system can act as a fish culture, plant cultivation and removal of ammoniacal nitrogen in water circulating system (aquaponic). Many researchers have studied the use of ABS to treat fish wastewater (Marques *et al.*, 2017) and marine fish wastewater treatment (Boxman *et al.*, 2018). However, the utilization of aquaponic system as a method to improve water quality in lake has not yet been well investigated. Therefore, the green technology of aquaponic system might produce effective system for ammoniacal nitrogen removal which provide a more efficient treatment of pollutant in lake water, and this emphasize the novelty of the current work. Aquaponic system used *Heliconia* sp. as a plant to uptake nutrient to clean the water. This method also used tilapia (*Oreochromis niloticus*) as a source of ammoniacal nitrogen to replace ammoniacal nitrogen from fertilizer in palm oil plantation. However, to the best of author's knowledge, not well documented research has been reported on the utilization of *Heliconia* sp. in removing ammoniacal nitrogen which was affected by photoperiod and temperature. Therefore, the effectiveness of ammoniacal nitrogen removal by ABS in lake water need to be investigated as it has potential to be an alternative method for photoperiod and temperature prepared by previous methods. Thus, this study aims to identify the characteristics of lake water to determine the removal efficiency of nutrients from lake water near palm oil plantation through ABS, and to optimize the different environmental effects of *Heliconia* sp. in terms of nutrient uptakes and potential of commercial value.

### 1.3 Objectives and research questions of study

The main aim of the current work is to investigate the efficiency of aquaponic biofilter for ammoniacal nitrogen removal using *Heliconia* sp. Plant. The specific objectives are as follows:

- i. To characterize the pollutants in *Tasik Teknologi* water.
  - a) What are the parameters and the mean value for concentration of ammoniacal nitrogen available in the lakes?
- ii. To determine the best operating parameters and efficiency of ABS for ammoniacal nitrogen removal.
  - a) What is the suitable temperature and photoperiod for Aquaponics Biofilter System?
  - b) How the most suitable operating condition for Aquaponics Biofilter System that will give the most effective ammoniacal nitrogen removal?
- iii. To analyze the nutrient composition on *Heliconia* sp. (Ornamental plant) after phytoremediation.
  - a) How is the value of ammoniacal nitrogen, nitrate and nitrite in root and leaves of the *Heliconia* sp.?

### 1.4 Scope of study

The present study focused on the removing of ammoniacal nitrogen from the lake which is located in Universiti Tun Hussein Onn Malaysia (UTHM) that surrounds the area of palm oil plantation. Lake water samples were obtained from the Faculty of Civil and Environmental Engineering Lake, *Tasik Teknologi* (1.8586° N, 103.0856° E) from 8.00 to 10.00 a.m. The samples of lake water were analyzed for biological oxygen demand (BOD), chemical oxygen demand (COD), total suspended solid (TSS), dissolved oxygen (DO) and ammoniacal nitrogen (AN). The aquaponic biofilter system was designed in a greenhouse scale using plant, *Heliconia* sp. and fish (tilapia, *Oreochromis niloticus*). The plants, of average 30 cm height and a tilapia with approximately 100 grams weight were



selected. The ABS was run for 10 days continuously. The best operating parameters (temperature and photoperiod) for aquaponic as biofilter system were determined based on the effectiveness of ammoniacal nitrogen removal.

#### 1.4 Hypotheses of the study

Several hypotheses can be derived through observation on an operation using Aquaponic Biofilter System (ABS) reactor as follows:

- a) If the runoff fertilizers (eutrophication) into the lake water that came from the palm oil plantation are high, then the lake water will be rich in nutrient contents such as nitrogen (ammoniacal nitrogen) and phosphorus.
- b) If the relationship between different environmental culture conditions of *Heliconia* sp. and specific growth rates exist, then the growth of *Heliconia* sp. will be different in each environmental culture condition.
- c) If the relationship between environmental factors such as temperature and photoperiod and specific growth of *Heliconia* sp. exist, then different factors will affect the growth rates and efficiency of ABS on nutrients removal.
- d) If the different environmental culture conditions of *Heliconia* sp. influence the phytoremediation of pollutants from lake water, then different culture conditions will result in different ABS efficiencies.
- e) If the different temperature and photoperiod influence the uptake of pollutant (ammoniacal nitrogen, nitrate and nitrite) in leaves, stems and roots of *Heliconia* sp., then there are one suitable temperature and photoperiod that contains high concentration ammoniacal nitrogen, nitrate and nitrite in leaves, stems and roots.

#### 1.6 Significance of study

The presence of ammoniacal nitrogen in water can pose both threat and benefits. The excessive, ammoniacal nitrogen can result in the death of aquatic organisms. However, at its safe level, ammoniacal nitrogen can reduce algae bloom, create clean and non-greenish water, avoid aquatic organism from dead, and remove pungent smell from water.

Optimization of parameters for ABS can determine if *Heliconia* sp. is suitable for aquaponic in Malaysia. The best concentration of carbon dioxide, water temperature and photoperiod for plant to grow well and healthy can also obtained from the optimization study. Furthermore, this research has identified the performance of *Heliconia* sp. as an ammoniacal nitrogen removal in the investigated lake.

The application of ABS can offer many benefits too. Its application in the investigated lake can help to reduce the level of excessive ammoniacal nitrogen that is coming from palm oil plantation. ABS can also compete with other useless plant like weed and water lily from growing.

## **1.7 Thesis structure**

This thesis consists of five chapters. Chapter 1 presents general information on palm oil plantation, ABS and plant potential in removal of nutrients. It is presented in few segments started with the background of study, followed by the problem statement, objectives of study, scope of study, significance of study, and closed with thesis structure.

Chapter 2 contains the review of different topics related to this research. This chapter presents information on the lake water in Malaysia, conventional methods for ammoniacal nitrogen removal, phytoremediation process, selection of fish (*Oreochromis niloticus*), selection of plant (*Heliconia* sp.), plant growth at different factors such as temperature and photoperiod of ABS and the commercial value of product of ABS. Major references are critically reviewed to give a clear view on what is expected from the current effort.

Chapter 3 describes in detail the methodology of this research. The first part describes the sampling process of lake water and the analysis conducted on several



## REFERENCES

- Abdel--Raouf, N., Al--Homaidan, A. A., & Ibraheem, I. B. M. (2012). Microalgae and wastewater treatment. *Saudi journal of biological sciences*, 19(3), 257--275.
- Abdullah, N., & Sulaiman, F. (2013). The palm oil wastes in Malaysia. In *Biomass Now--Sustainable Growth and Use*. InTech
- Abdullah, R., & Wahid, M. B. (2010). World palm oil supply, demand, price and prospects: focus on Malaysian and Indonesian palm oil industry. *Malaysian Palm Oil Board Press, Malaysia*.
- Abe, K., Ozaki, Y., & Mizuta, K. (1999). Evaluation of useful plants for the treatment of polluted pond water with low N and P concentrations. *Soil science and plant nutrition*, 45(2), 409-417..
- Akhir, N. I. M., Kusin, F. M., Mohamat-Yusuff, F., Awang, M., & Hamka, F. (2014). Nitrous oxide emission from different land use changes associated with oil palm plantation. *Journal of Agriculture and Environmental Sciences*, 3(2), 421-431..
- Al--Badaai, F., Shuhaimi--Othman, M., & Gasim, M. B. (2013). Water quality assessment of the Semenyih river, Selangor, Malaysia. *Journal of chemistry*, 2013.
- Ansar, M. (2018). Sustainable integrated farming system: A solution for national food security and sovereignty. In *IOP Conference Series: Earth and Environmental Science* (Vol. 157, No. 1, p. 012061). IOP Publishing.
- Anandha Varun, R., & Kalpana, S. (2015). Performance analysis of nutrient removal in pond water using Water Hyacinth and Azolla with papaya stem. *Int Res J Eng Technol*, 2(1), 444--448.
- Anonymus, (2014). Ammoniacal nitrogen in groundwater, runoff and stream.

- APHA (2005). Standard Methods for the Examination of Water and Wastewater. 21st Edition, American Public Health Association/American Water Works Association/Water Environment Federation, Washington DC.
- ASM. (2010), Lake Briefs Synthesis Report. Managing Lakes and Their Basins for Sustainable Use in Malaysia (Lake Briefs Report Series 1). Kuala Lumpur: Akademi Sains Malaysia.
- ASM, NAHRIM. (2009). Pelan Strategi: Strategi Pembangunan dan Pengurusan Lestari bagi - Tasik dan Empangan Air di Malaysia Jilid 1: Laporan Utama.
- Asharuddin, M. S. Zayadi, N. Rasit, W. & Othman, N. (2016) Water Quality Characteristics of Sembrong Dam Reservoir, Johor, Malaysia, *IOP Conf. Ser. Mater. Sci. Eng.*, vol. 136, no. 1, pp. 6–12, 2016.
- Bah, A., Husni, M. H. A., Teh, C. B. S., Rafii, M. Y., Syed Omar, S. R., & Ahmed, O. H. (2014). Reducing runoff loss of applied nutrients in palm oil cultivation using controlled-release fertilizers. *Advances in Agriculture*, 2014.
- Bernstein (2013). “This is an environmentally friendly, natural food-growing method that harnesses the best attributes”. retrieved from
- Bindu, G. H. (2015). Impact Of Toxic Metals, Minerals, Solvents, E-Waste And Plastics Leading To Environmental Pollution. *Journal of Chemical and Pharmaceutical Sciences* ISSN, 974, 2115.
- Blamey, M. & Grey-Wilson, C. (1989). Flora of Britain and Northern Europe. ISBN 0-340-40170-2.
- Bohutskyi, P., Liu, K., Nasr, L. K., Byers, N., Rosenberg, J. N., Oyler, G. A., & Bouwer, E. J. (2015). Bioprospecting of microalgae for integrated biomass production and phytoremediation of unsterilized wastewater and anaerobic digestion centrate. *Applied microbiology and biotechnology*, 99(14), 6139-6154.

- Bostock, J., McAndrew, B., Richards, R., Jauncey, K., Telfer, T., Lorenzen, K., & Corner, R. (2010). Aquaculture: global status and trends. *Philosophical Transactions of the Royal Society of London B: Biological Sciences*, 365(1554), 2897–2912.
- Boxman, S. E., Nystrom, M., Ergas, S. J., Main, K. L., & Trotz, M. A. (2018). Evaluation of water treatment capacity, nutrient cycling, and biomass production in a marine aquaponic system. *Ecological Engineering*, 120, 299–310.
- Braadbaart, O. & Braadbaart, F. (1997). Policing the Urban Pumping Race: Industrial Groundwater Overexploitation in Indonesia. *World Development*, 25(2). (pp 199–210).
- Breisha, G.Z., Winter, J. (2010), Bio—removal of nitrogen from wastewaters—a review, *Am. J. Sci.* 6 508–528.
- Brix, H., Koottatep, T., & Laugesen, C. H. (2007). Wastewater treatment in tsunami affected areas of Thailand by constructed wetlands. *Water Science and Technology*, 56(3), 69–74.
- Brooks, M.A., (1999), Breakpoint Chlorination as an Alternative Means of Ammoniacal nitrogen— Nitrogen Removal at a Water Reclamation Plant, M.S. Thesis in Environmental Sciences and Engineering, Virginia Polytechnic Institute and State University.
- Buzby, K. M., & Lin, L. S. (2014). Scaling aquaponic systems: Balancing plant uptake with fish output. *Aquacultural engineering*, 63, 39–44.
- Capodaglio, A.G., Hlavínek, P., Raboni, M., (2015), Physico—chemical technologies for nitrogen removal from wastewaters: a review, *Rev. Ambient. Água* 10 481–498.
- Catley, J. L., & Brooking, I. R. (1996). Temperature and Light Influence Growth and Flower Production in Heliconia Golden Torch'. *HortScience*, 31(2), 213–217.
- Chen, X., Wang, Z., Tang, S., Zhao, Y., & Zhao, J. (2017). Genome-wide mapping of DNA methylation in Nile Tilapia. *Hydrobiologia*, 791(1), 247–257.

- Chen, C., Zhao, T., Liu, R., & Luo, L. (2017). Performance of five plant species in removal of nitrogen and phosphorus from an experimental phytoremediation system in the Ningxia irrigation area. *Environmental monitoring and assessment*, 189(10), 497.
- Chen, Y., Wu, Y., Liu, C., Guo, L., Nie, J., Chen, Y., & Qiu, T. (2018). Low–temperature conversion of ammoniacal nitrogen to nitrogen in water with ozone over composite metal oxide catalyst. *Journal of Environmental Sciences*, 66, 265–273.
- Chooklin, S, Kaewsichan, L. & Kaewsrichan, J. (2013). Potential Use of Palm oil Sap on Lactic Acid Production and Product Adsorption on Dowex™ 66 Resin as Adsorbent. *Asia Pacific Journal ChemicalEngineering*, 8(1): 23 – 31.
- Comte, I., Colin, F., Whalen, J. K., Grünberger, O., & Caliman, J. P. (2012). Agricultural practices in palm oil plantations and their impact on hydrological changes, nutrient fluxes and water quality in Indonesia: a review. In *Advances in Agronomy* (Vol. 116, pp. 71–124). Academic Press.
- Correa, C. Z., Prates, K. V. M. C., de Oliveira, E. F., Lopes, D. D., & Barana, A. C. (2018). Nitrification/denitrification of real municipal wastewater in an intermittently aerated structured bed reactor. *Journal of Water Process Engineering*, 23, 134–141.
- Constable, M. Charlton, M. Jensen, F. McDonald, K. Craig, G. & Taylor, K. W. (2003) “An ecological risk assessment of ammoniacal nitrogen in the aquatic environment,” *Human and Ecological Risk Assessment*, vol. 9, no. 2, pp. 527–548.
- Criley, R.A. and T.K. Broschat. (1992). *Heliconia*: Botany and horticulture of a new floral crop. *Hort. Rev.* 14:1–55.
- Culp, R.L., Culp, G.L., (1971) *Advanced wastewater treatment*, Van Nostrand Reinhold, The Netherlands.
- Daud, N. N. N., & Syazwani, A. A. (2016). Preliminary assessment of lakes water quality status at campus area in selangor, Malaysia.
- Dediu, L., Cristea, V., & Xiaoshuan, Z. (2012). Waste production and valorization in an integrated aquaponic system with bester and lettuce. *African Journal of Biotechnology*, 11(9), 2349–2358.

- Del-Pozo, J., Mishra, N., Kabuusu, R., Cheetham, S., Eldar, A., Bacharach, E., Lipkin, W.I., Ferguson, H.W., (2017). Syncytial hepatitis of tilapia (*Oreochromis niloticus* L.) is associated with Orthomyxovirus–like virions in hepatocytes. *Vet. Pathol.* 54, 164–170.
- Delkash, M., Al-Faraj, F. A., & Scholz, M. (2018). Impacts of anthropogenic land use changes on nutrient concentrations in surface waterbodies: a review. *CLEAN–Soil, Air, Water*, 46(5), 1800051.
- De Silva, S. S. (2012). Aquaculture: a newly emergent food production sector—and perspectives of its impacts on biodiversity and conservation. *Biodiversity and conservation*, 21(12), 3187–3220.
- Ding, S., Bao, P., Wang, B., Zhang, Q., & Peng, Y. (2018). Long–term stable simultaneous partial nitrification, anammox and denitrification (SNAD) process treating real domestic sewage using suspended activated sludge. *Chemical Engineering Journal*, 339, 180–188.
- Dressing, S. A., Meals, D. W., Harcum, J. B., Spooner, J., Stribling, J. B., Richards, R. P., & O'Donnell, J. G. (2016). *Monitoring and Evaluating Nonpoint Source Watershed Projects*. EPA 841–R–16–010. US Environmental Protection Agency, Washington, DC. Accessed December 20, 2016. <https://www.epa.gov/nps/monitoring--and--evaluating--nonpoint--source--watershed--projects>.
- Effendi, H., Utomo, B. A., & Pratiwi, N. T. (2018). Ammoniacal nitrogen and orthophosphate removal of tilapia cultivation wastewater with *Vetiveria zizanioides*. *Journal of King Saud University–Science*.
- El–Khateeb, M. A., & El–Bahrawy, A. Z. (2013). Extensive post treatment using constructed wetland. *Life Science Journal*, 10(2), 560–568.
- Endut, A., Jusoh, A., Ali, N., Nik, W. W., & Hassan, A. (2010). A study on the optimal hydraulic loading rate and plant ratios in recirculation aquaponic system. *Bioresource technology*, 101(5), 1511–1517.

- Environmental Protection Agency (2005). *Evaluation of Phytoremediation for Management of Chlorinated Solvents in Soil and Groundwater*. United States: Environmental Protection Agency.
- Estim, A., Saufie, S., & Mustafa, S.(2018) *Water quality remediation using aquaponics sub--systems as biological and mechanical filters in aquaculture*, Journal of Water Process Engineering 2018.
- FAO, 2017a. Global aquaculture production.
- Fazal, S., Zhang, B., & Mehmood, Q. (2015). Biological treatment of combined industrial wastewater. *Ecological Engineering*, 84, 551-558.
- Fang, Y. Y., Yang, X. E., Chang, H. Q., Pu, P. M., Ding, X. F., & Rengel, Z. (2007). Phytoremediation of nitrogen--polluted water using water hyacinth. *Journal of plant nutrition*, 30(11), 1753--1765.
- Feng, Q., Liu, G., Meng, L., Fu, E., Zhang, H. & Zhang, K. (2008). Land Subsidence Induced by Groundwater Extraction and Building Damage Level Assessment -- A Case Study of Datun, China. *Journal of China University of Mining and Technology* 18, 556 -- 560.
- Fitzsimmons, K., (2000). Future trends of tilapia aquaculture in the Americas, in: Costa--Pierce, B.A., Rakocy, J.E. (Eds.), *Tilapia Aquaculture in the Americas*, Vol. 2. The World Aquaculture Society, Baton Rouge, Louisiana, United States, pp 252--264.
- FOMCA, WECAM & Malaysian Water Forum. (2009). No Recharge, No Discharge, Sime Darby Will Be Charge. Research Report on Groundwater Project in Batang Padang, Perak.
- Food and Agriculture Organisation, FAO (2005). Malaysia -- FAO's Information System on Water and Agriculture. Encyclopedia of Earth.Cleveland (Washington, D.C.:



- Environmental Information Coalition, National Council for Science and the Environment), United Nations. Retrieved on October 2, 2012 from
- Fox, L. J., Struik, P. C., Appleton, B. L., & Rule, J. H. (2008). Nitrogen phytoremediation by water hyacinth (*Eichhornia crassipes* (Mart.) Solms). *Water, Air, and Soil Pollution*, 194(1-4), 199-207.
- Fry, J. P., Love, D. C., MacDonald, G. K., West, P. C., Engstrom, P. M., Nachman, K. E., & Lawrence, R. S. (2016). Environmental health impacts of feeding crops to farmed fish. *Environment international*, 91, 201–214.
- Gasim, M. B., Toriman, M. E., Muftah, S., Barggig, A., Aziz, N. A. A., Azaman, F., & Muhamad, H. (2015). Water quality degradation of Cempaka Lake, Bangi, Selangor, Malaysia as an impact of excessive *E. coli* and nutrient concentrations. *Malaysian Journal of Analytical Sciences*, 19(6), 1391–1404.
- Geng, Y., Baumann, F., Song, C., Zhang, M., Shi, Y., Kühn, P., & He, J. S. (2017). Increasing temperature reduces the coupling between available nitrogen and phosphorus in soils of Chinese grasslands. *Scientific Reports*, 7, 43524.
- Giustinianovich, E.A., Campos, J., Roeckel, M.D., (2016), The presence of organic matter during autotrophic nitrogen removal: problem or opportunity? Sep. Purif. Technol. 166 102–108.
- Goh, K. J., Teo, C. B., Chew, P. S., & Chiu, S. B. (1999). Fertiliser management in palm oil: Agronomic principles and field practices. *Fertiliser management for palm oil plantations*, 20–21.
- Goddek, S. (2017). *Opportunities and challenges of multi-loop aquaponic systems* (Doctoral dissertation, Wageningen University).
- Gomes, H.I., (2012) Phytoremediation for bioenergy: challenges and opportunities, *Environ. Technol. Rev.* 1 59–66.
- Gupta, V.K., Sadegh, H., Yari, M. Shahryari Ghoshekandi, R., Maazinejad, B., Chahardori, M., (2015), Removal of ammonium ions from wastewater: a short

review in development of efficient methods, *Glob. J. Environ. Sci. Manage.* 1 149–158.

Ha, T. M., Krisantini, S., & Johnston, M. E. (2013). The effect of photoperiod and temperature on flowering of *Pycnosorus thompsonianus*. *Asian Journal of Agriculture and Food Science (ISSN: 2321–1571)*, 1(05).

Hairston N. G. & Fussmann, G. F. (2014) *Lake Ecosystems, Els.*

Harris, L. M., Goldin, J. A., & Sneddon, C. (Eds.). (2015). *Contemporary water governance in the global south: Scarcity, marketization and participation*. Routledge.

Hamid, S., Bae, S., & Lee, W. (2018). Novel bimetallic catalyst supported by red mud for enhanced nitrate reduction. *Chemical Engineering Journal*, 348, 877–887.

Haseena, P. V., Padmavathy, K. S., Krishnan, P. R., & Madhu, G. (2016). Adsorption of Ammonium Nitrogen from Aqueous Systems Using Chitosan–Bentonite Film Composite. *Procedia Technology*, 24, 733–740.

Hart, M. R., Quin, B. F., & Nguyen, M. (2004). Phosphorus runoff from agricultural land and direct fertilizer effects. *Journal of environmental quality*, 33(6), 1954–1972.

Hayes, N. M., Deemer, B. R., Corman, J. R., Razavi, N. R., & Strock, K. E. (2017). Key differences between lakes and reservoirs modify climate signals: A case for a new conceptual model. *Limnology and Oceanography Letters*, 2(2), 47–62.

Hidzrami S.A. (2010), Bukit Merah, Perak Lake Brief. Lakes and Their Basins for Sustainable Use in Malaysia (Lake Briefs Report Series 1). Kuala Lumpur: Akademi Sains Malaysia.

Holm, B., Heinsoo, K., (2013). Municipal wastewater application to Short Rotation Coppice of willow: Treatment efficiency and clone response in Estonian case study. *Biomass Bioenergy* 57, 126–135.

Hooda, P. S., Edwards, A. C., Anderson, H. A., & Miller, A. (2000). A review of water quality concerns in livestock farming areas. *Science of the total environment*, 250(1–3), 143–167.



- Hu, Z., Lee, J. W., Chandran, K., Kim, S., Brotto, A. C., & Khanal, S. K. (2015). Effect of plant species on nitrogen recovery in aquaponics. *Bioresource technology*, 188, 92–98.
- Huang, H., Xiao, D., Zhang, Q., & Ding, L. (2014). Removal of ammoniacal nitrogen from landfill leachate by struvite precipitation with the use of low–cost phosphate and magnesium sources. *Journal of environmental management*, 145, 191–198.
- ILEC. (2007) Integrated Lake Basin Management: An Introduction. Kusatsu, Japan: International Lake Environment Committee Foundation.
- Ilyas, H., Masih, I., (2017) The performance of the intensified constructed wetlands for organic matter and nitrogen removal: a review, *J. Environ. Manage.* 198 372–383.
- International Water Association, (2000). *Constructed Wetlands for Pollution Control. Processes, Performance, Design and Operation*. IWA Publishing, London.
- Inhaber, H. (1975), An approach to a water quality index for Canada, *Water Res.*, vol. 9, no. 9, pp. 821–833, 1975.
- Jasrotia, S., Kansal, A., & Mehra, A. (2017). Performance of aquatic plant species for phytoremediation of arsenic–contaminated water. *Applied Water Science*, 7(2), 889–896.
- Garrido, J. M., Guerrero, L., Méndez, R., & Lema, J. M. (1998). Nitrification of waste waters from fish-meal factories. *Water S. A.*, 24(3), 245–251..
- Joesting, H. M., Blaylock, R., Biber, P., & Ray, A. (2016). The use of marine aquaculture solid waste for nursery production of the salt marsh plants *Spartina alterniflora* and *Juncus roemerianus*. *Aquaculture reports*, 3, 108–114.
- Jorgensen, S. E., & Williams, W. D. (2001). Water quality: The impact of eutrophication. *UN Environment Programme*.
- Khan, F. A., & Ansari, A. A. (2005). Eutrophication: an ecological vision. *The botanical review*, 71(4), 449–482.

- Kianirad, E. Bedoya, D. Mcgarvey, K. and Novotny, V. (2006) Review of Watershed Ecological Models Center for Urban Environmental Studies Review of Watershed Ecological Models Student Investigators, 2006.
- Khan, F. A., & Ansari, A. A. (2005). Eutrophication: an ecological vision. *The botanical review*, 71(4), 449-482.
- Kusin, F. M., Akhir, N. I. M., Mohamat--Yusuff, F., & Awang, M. (2015). The impact of nitrogen fertilizer use on greenhouse gas emissions in an palm oil plantation associated with land use change. *Atmósfera*, 28(4), 243–250
- Konnerup, D., Koottatep, T., Brix, H., (2009). Treatment of domestic wastewater in tropical, subsurface flow constructed wetlands planted with Canna and Heliconia. *Ecol. Eng.* 35, 248e257. <http://dx.doi.org/10.1016/j.ecoleng.2008.04.018>
- Kowalik, P.J., Randerson, P.F., (1994). Nitrogen and phosphorus removal by willow stands irrigated with municipal waste waterda review of the Polish experience. *Biomass Bioenergy* 6, 133e139. [http://dx.doi.org/10.1016/0961-9534\(94\)90092-2](http://dx.doi.org/10.1016/0961-9534(94)90092-2).
- Lam, S. S., Ma, N. L., Jusoh, A., & Ambak, M. A. (2015). Biological nutrient removal by recirculating aquaponic system: Optimization of the dimension ratio between the hydroponic & rearing tank components. *International Biodeterioration & Biodegradation*, 102, 107–115.
- Landwehr, J. M. & Deininger, R.A. (1976) A comparison of several water quality indexes., *Water Pollut. Control Fed.*, pp. 954–958.
- Li, L., Dong, Y., Qian, G., Hu, X., & Ye, L. (2018). Performance and microbial community analysis of bio–electrocoagulation on simultaneous nitrification and denitrification in submerged membrane bioreactor at limited dissolved oxygen. *Bioresource technology*, 258, 168–176.
- Li, R., Feng, C., Hu, W., Xi, B., Chen, N., Zhao, B., & Pu, J. (2016). Woodchip–sulfur based heterotrophic and autotrophic denitrification (WSHAD) process for nitrate

contaminated water remediation. *Water research*, 89, 171–179.

Lim CE, Webster CD, Li MH. (2006). Feeding practices. In: Lim CE, Webster CD, editors. *Tilapia biology, culture, and nutrition*. Binghamton: The Haworth Press; p. 547–559.

Lima, R.E.M.; Farias, F.C.; Bezerra, M.A.; Bezerra, F.C. (2016). Parâmetros biométricos e fisiológicos de *Heliconia bihai* cultivada em região litorânea sob diferentes níveis de radiação solar. *Ornamental Horticulture*, v.22, n.1, p.50–57. DOI:

Lu, Q., He, Z. L., Graetz, D. A., Stoffella, P. J., & Yang, X. (2010). Phytoremediation to remove nutrients and improve eutrophic stormwaters using water lettuce (*Pistia stratiotes* L.). *Environmental Science and Pollution Research*, 17(1), 84-96.

Luo, X., Yan, Q., Wang, C., Luo, C., Zhou, N., & Jian, C. (2015). Treatment of ammoniacal nitrogen wastewater in low concentration by two-stage ozonization. *International journal of environmental research and public health*, 12(9), 11975–11987.

Lynch, A. J., Cooke, S. J., Deines, A. M., Bower, S. D., Bunnell, D. B., Cowx, I. G., & Rogers, M. W. (2016). The social, economic, and environmental importance of inland fish and fisheries. *Environmental Reviews*, 24(2), 115–121.

Ma, B., Wang, S., Cao, S., Miao, Y., Jia, F., Du, R., Peng, Y., (2016), Biological nitrogen removal from sewage via anammox: recent advances, *Bioresour. Technol.* 200 (2016) 981–990.

Madaki, Y. S., & Seng, L. (2013). Palm oil mill effluent (POME) from Malaysia palm oil mills: waste or resource. *International Journal of Science, Environment and Technology*, 2(6), 1138–1155.

- Mahieu, A. (2015). *Fish--farming in South Africa: A study of the market environment and the suitable species* (Doctoral dissertation, Stellenbosch: Stellenbosch University).
- Malaysian Palm Oil Board, MPOB (2016). Overview of the Malaysia Palm oil Industry by Economics & Industry Development Division MPOB on January 2017 retrieved from [http://bepi.mpob.gov.my/images/overview/Overview\\_of\\_Industry\\_2016.pdf](http://bepi.mpob.gov.my/images/overview/Overview_of_Industry_2016.pdf)
- Mamat NZ, Shaari MI, Wahab NAAA (2016) The Production of Catfish and Vegetables in an Aquaponic System. *Fish Aquac J* 7: 181.doi:10.4172/2150--3508.1000181.
- Manarangi, A., W.S. Sakai, C. Gerken, M. Crowell, G. Nielson, and R. Short. (1988). Growth and flowering of *Heliconia psittacorum* cv. Parrot. *Hawaii J. Hawaii Pacific Agr.* 1:1--3.
- Marques, B., Calado, R., & Lillebø, A. I. (2017). New species for the biomitigation of a super--intensive marine fish farm effluent: Combined use of polychaete--assisted sand filters and halophyte aquaponics. *Science of the Total Environment*, 599, 1922--1928.
- Mayo, A. W., & Hanai, E. E. (2017). Modeling phytoremediation of nitrogen-polluted water using water hyacinth (*Eichhornia crassipes*). *Physics and Chemistry of the Earth, Parts A/B/C*, 100, 170-180.
- Mekonnen MM, Hoekstra AY (2010) The green, blue and grey water footprint of crops and derived crop products. (UNESCO--IHE, Delft, The Netherlands) Value of Water Research Report Series No. 47.
- Mesquita, M. C., Albuquerque, A., Amaral, L., & Nogueira, R. (2017). Seasonal variation of nutrient removal in a full--scale horizontal constructed wetland. *Energy Procedia*, 136, 225--232.
- Melati Mohd Ariff, (2010). Malaysia: No Water, No Life. *Bernama*, Retrieved on 15 March 2011.

Méndez–Mendoza, A. S., Bello–Mendoza, R., Herrera–López, D., Mejía–González, G., & Calixto–Romo, A. (2015). Performance of constructed wetlands with ornamental plants in the treatment of domestic wastewater under the tropical climate of South Mexico. *Water Practice and Technology*, 10(1), 110–123.

Merino, G., Barange, M., Blanchard, J. L., Harle, J., Holmes, R., Allen, I., & Jennings, S. (2012). Can marine fisheries and aquaculture meet fish demand from a growing human population in a changing climate, *Global Environmental Change*, 22(4), 795–806.

Mishra, S., & Maiti, A. (2017). The efficiency of *Eichhornia crassipes* in the removal of organic and inorganic pollutants from wastewater: a review. *Environmental science and pollution research*, 24(9), 7921-7937.

Mokhtari-Hosseini, Z. B., Ehsan, K., Reza, T., & Toktam, S. Z. (2016). Optimization of ammonia removal by natural zeolite from aqueous solution using response surface methodology. *Hemijiska Industrija*, 70(1), 21.

Morand, P., Robin, P., Escande, A., Picot, B., Pourcher, A. M., Jiangping, Q., & Oudart, D. (2011). Biomass production and water purification from fresh liquid manure—Use of vermiculture, macrophytes ponds and constructed wetlands to recover nutrients and recycle water for flushing in pig housing. *Procedia Environmental Sciences*, 9, 130-139.

Nadzir, N. S. C., Abdullah, M. Z., & Sulaiman, F. R. (2019). Surface water quality in palm oil plantation. *Malaysian Journal of Fundamental and Applied Sciences*, 15(1), 85-87.

Nancharaiah, Y.V., Venkata Mohan, S., Lens, P.N.L, (2016), Recent advances in nutrient removal and recovery in biological and bioelectrochemical systems, *Bioresour. Technol.* 215 173–185.

National Water Quality Standards for Malaysia. (2006). Retrieved from [http://www.wepa-db.net/policies/law/malaysia/eq\\_surface.htm](http://www.wepa-db.net/policies/law/malaysia/eq_surface.htm).

Naidoo, S., & Olaniran, A. O. (2013). Treated wastewater effluent as a source of microbial pollution of surface water resources. *International journal of environmental research and public health*, 11(1), 249–270.

- NMSU DACC WUTAP, (2007) New Mexico Wastewater Systems Operator Certification Study Manual, Chapter 13: Nitrogen Removal, Version 1.1, pp. 162–172.
- Parra, C. A. M. (2016). Treatment of landfill leachate by polyculture constructed wetlands planted with native plants. *Ingeniería y competitividad: revista científica y tecnológica*, (2), 183–192.
- Peña–Salamanca, E. J., Madera–Parra, C. A., Sánchez, J. M., & Medina–Vásquez, J. (2013). Bioprospecting of native plants for their use in bioremediation process: *Heliconia psittacorum* case (heliconiaceae). *Revista de la Academia Colombiana de Ciencias Exactas, Físicas y Naturales*, 37(145), 469–481.
- Pirker, J., Mosnier, A., Kraxner, F., Havlík, P., & Obersteiner, M. (2016). What are the limits to palm oil expansion, *Global Environmental Change*, 40, 73–81.
- Popp, J., Váradi, L., Békefi, E., Péteri, A., Gyalog, G., Lakner, Z., & Oláh, J. (2018). Evolution of integrated open aquaculture systems in Hungary: results from a case study. *Sustainability*, 10(1), 177.
- Provolo, G., Perazzolo, F., Mattachini, G., Finzi, A., Naldi, E., & Riva, E. (2017). Nitrogen removal from digested slurries using a simplified ammoniacal nitrogen stripping technique. *Waste Management*, 69, 154–161.
- Ojoawo, S. O., Udayakumar, G., & Naik, P. (2015). Phytoremediation of phosphorus and nitrogen with *Canna x generalis* reeds in domestic wastewater through NMAMIT constructed wetland. *Aquatic Procedia*, 4, 349–356.
- Orentlicher, M. (2012) Overview of nitrogen removal technologies and application/use of associated end products, Proceeding of Got Manure Tradeshow and Conference, Cornell University, New York, 2012.
- Ramadas, K. (2009). Report on Governance Sector: Position Paper on Governance of Lakes in Malaysia (2008). *Strategy Plan: Strategies for the Sustainable Development and Management of Lakes and Reservoirs in Malaysia*, 2.
- Rahmani, A.R., Mahvi, A.H., (2006) Use of ion exchange for removal of ammonium: a biological regeneration of zeolite, *Glob. NEST J.* 8 146–150.



- Reantaso, M.B., (2017). Tilapialake virus (TiLV) caused by an orthomyxo-like virus (Family Orthomyxoviridae) threatening cultured and wild stocks of tilapia. *FAO Aquaculture Newsletter* 57, 9-11.
- Reddy, K.R., D'Angelo, E.M., (1990), Biomass yield and nutrient removal by water hyacinth (*Eichhornia crassipes*) as influenced by harvesting frequency, *Biomass* 21 27–42.
- Rice, E.W., Bridgewater, L., Association, A.P.H.A, (2012). *Standard Methods for the Examination of Water and Wastewater*, twenty–second ed. American Public Health Association, Washington, DC.
- Robert M. D., David M. C. and Martin W. B. (1997). Ammoniacal nitrogen in Fish Ponds. Southern Regional Aquaculture Center (SRAC). Publication No. 463.
- Robertson, G.P., Groffman, P.M., (2015) Nitrogen transformations, in: E.A. Paul (Ed.), *Soil Microbiology, Ecology and Biochemistry*, 4th edition, Academic Press, Burlington, Massachusetts, USA, 2015, pp. 421–446.
- Rysgaard, S., Risgaard–Petersen, N., Nielsen, L. P., & Revsbech, N. P. (1993). Nitrification and denitrification in lake and estuarine sediments measured by the  $^{15}\text{N}$  dilution technique and isotope pairing. *Applied and Environmental Microbiology*, 59(7), 2093–2098.
- Saha, S., Monroe, A., & Day, M. R. (2016). Growth, yield, plant quality and nutrition of basil (*Ocimum basilicum* L.) under soilless agricultural systems. *Annals of Agricultural Sciences*, 61(2), 181–186.
- Saimy, I. S., & Yusof, N. A. M. (2013). The need for better water policy and governance in Malaysia. *Procedia–Social and Behavioral Sciences*, 81, 371–375.
- Saleh, H. M. (2012). Water hyacinth for phytoremediation of radioactive waste simulate contaminated with cesium and cobalt radionuclides. *Nuclear Engineering and Design*, 242, 425–432.
- Salameh, E., Harahsheh, S. (2011) Eutrophication Processes in Arid Climates. In Ansari AA, Gill SS, Lanza GR, Rast W (eds.) *Eutrophication: Causes, Consequences and Control*. New York: Springer.

- Sampson, B. (2013). Wastewater sampling. Athens, Georgia: Region 4 U.S. environmental protection agency Science and ecosystem support Division Athens, Georgia.
- Savci, S. (2012). Investigation of effect of chemical fertilizers on environment. *Apacbee Procedia*, 1, 287–292.
- Sharip, Z., Zaki, A. T., Shapai, M., Suratman, S., & Shaaban, A. J. (2014). Lakes of Malaysia: Water quality, eutrophication and management. *Lakes & Reservoirs: Research & Management*, 19(2), 130–141.
- Silva, L., Valdés–Lozano, D., Escalante, E., & Gasca–Leyva, E. (2018). Dynamic root floating technique: An option to reduce electric power consumption in aquaponic systems. *Journal of Cleaner Production*, 183, 132–142.
- Sim, C. H., Yusoff, M. K., Shutes, B., Ho, S. C., & Mansor, M. (2008). Nutrient removal in a pilot and full scale constructed wetland, Putrajaya city, Malaysia. *Journal of Environmental Management*, 88(2), 307–317.
- Smith, D. G. (1990) A better water quality indexing system for rivers and streams, *Water Res.*, vol. 24, no. 10, pp. 1237–1244.
- Sun, X., Du, L., Hou, Y., Cheng, S., Zhang, X., & Liu, B. (2018). Endogenous influences on anammox and sulfocompound–oxidizing autotrophic denitrification coupling system (A/SAD) and dynamic operating strategy. *Bioresource technology*.
- SPU. (2014), Sarawak Integrated Water Resources Management Master Plan Study. State Planning Unit Sarawak;2008.
- Szatkowska, B., Płaza, E., Trela, J. (2008), Partial nitrification/Anammox and CANON – nitrogen removal systems followed by conductivity measurements, Proceedings of Polish–Swedish seminars. Conference paper 13 109–117.
- Teiri, H., Pourzamani, H., & Hajizadeh, Y. (2018). Phytoremediation of VOCs from indoor air by ornamental potted plants: a pilot study using a palm species under the controlled environment. *Chemosphere*, 197, 375–381.



- Turcios, A. E., & Papenbrock, J. (2014). Sustainable treatment of aquaculture effluents—what can we learn from the past for the future, *Sustainability*, 6(2), 836–856.
- Turkmen, G., & Guner, Y. (2010). Aquaponic (Integrating Fish and Plant Culture) Systems.
- UNESCO (2012). “Considering the agricultural use of water alone, there will be an increase in global consumption demand of 19% by 2050”. Retrieved from
- United Nations (2013), Department of Economic and Social Affairs, Population Division. World Population Prospects: The 2012 Revision, Volume II, Demographic Profiles.
- USEPA (1999). Update of ambient water quality criteria for ammoniacal nitrogen — technical version—1999. EPA–823–F–99–024. USEPA, Washington DC, USA.
- Valipour, A., Hamnabard, N., Woo, K., Ahn, Y., (2014) Performance of high–rate constructed phytoremediation process with attached growth for domestic wastewater treatment: effect of high TDS and Cu, *J. Environ. Manage.* 145 1–8.
- Valipour, A., Azizi, S., Raman, V. K., Jamshidi, S., & Hamnabard, N. (2014). The Comparative Evaluation of the Performance of Two Phytoremediation Systems for Domestic Wastewater Treatment. *Journal of environmental science & engineering*, 56(3), 319–326
- Vairappan, C.S. & Yen A.M. (2008). Palm oil mill effluent (POME) cultured marine microalgae a supplementary diet for rotifer culture. *J. Appl. Physics*, 20(5): 153–158.
- Vargas, L. E. P., Laurance, W. F., Clements, G. R., & Edwards, W. (2015). The impacts of palm oil agriculture on Colombia's biodiversity: what we know and still need to know. *Tropical Conservation Science*, 8(3), 828–845.
- Von Uexküll, H. R. (2007). Palm oil (*Elaeis guineensis* Jacq.). E. & SE Asia Program for the Potash & Phosphate Institute. *International Potash Institute, Singapore*.
- Wang, C., Liu, S., Xu, X., Zhang, C., Wang, D., & Yang, F. (2018). Achieving mainstream nitrogen removal through simultaneous partial nitrification, anammox

- and denitrification process in an integrated fixed film activated sludge reactor. *Chemosphere*, 203, 457–466.
- Wang, C. Y., Chang, C. Y., Chien, Y. H., & Lai, H. T. (2016). The performance of coupling membrane filtration in recirculating aquaponic system for tilapia culture. *International Biodeterioration & Biodegradation*, 107, 21–30.
- Witus, I. W., & Vun, L. W. (2016). Aquaculture in Malaysia: a short review on current policy and legislation. *Transactions on Science and Technology*, 3(1–2), 150–154.
- Wieben, C., Baker, R. J., & Nicholson, R. S. (2013). *Nutrient Concentrations in Surface Water and Groundwater, and Nitrate Source Identification Using Stable Isotope Analysis in the Barnegat Bay–Little Egg Harbor Watershed, New Jersey, 2010–11*. US Department of the Interior, US Geological Survey.
- Wongkiew, S., Popp, B. N., Kim, H. J., Khanal, S. K., (2017)., Fate of nitrogen in floating raft aquaponic systems using natural abundance nitrogen isotope compositions, *International Biodeterioration & Biodegradation* 125 (2017) 24–32.
- World Health Organisation, WHO (2011). World Water Day 2006. Resource Sheet Number 2. Retrieved 21 November 2011 from <http://www.worldwaterday.org/>.
- Wright, R. T. (2008). *Environmental Science; Towards a Sustainable Future*, 10th Ed. New Jersey, Pearson Prentice Hall.
- Wurts, W. A. (2003). Daily pH cycle and ammonia toxicity. *World Aquaculture*, 34(2), 20–21.
- Xu, Y., Ibrahim, I. M., & Harvey, P. J. (2016). The influence of photoperiod and light intensity on the growth and photosynthesis of *Dunaliella salina* (chlorophyta) CCAP 19/30. *Plant Physiology and Biochemistry*, 106, 305–315.
- Yacob, S., Hassan, M. A., Shirai, Y., Wakisaka, M., & Subash, S. (2005). Baseline study of methane emission from open digesting tanks of palm oil mill effluent treatment. *Chemosphere*, 59(11), 1575–1581.
- Yavari, S., Malakahmad, A. M. I. R. H. O. S. S. E. I. N., & Sapari, N. B. (2017). Phytoremediation of urea fertilizer factory wastewater by teak (*Tectona grandis*). *Environment Protection Engineering*, 43(4).

- Yavuzcan Yildiz, H., Robaina, L., Pirhonen, J., Mente, E., Domínguez, D., & Parisi, G. (2017). Fish Welfare in Aquaponic Systems: Its Relation to Water Quality with an Emphasis on Feed and Faeces—A Review. *Water*, 9(1), 13.
- Ye, L., Li, D., Zhang, J., & Zeng, H. (2018). Resuscitation of starved anaerobic ammonium oxidation sludge system: Impacts of repeated short-term starvation. *Bioresource technology*, 263, 458–466.
- Yue, X., Yu, G., Lu, Y., Liu, Z., Li, Q., Tang, J., & Liu, J. (2018). Effect of dissolved oxygen on nitrogen removal and the microbial community of the completely autotrophic nitrogen removal over nitrite process in a submerged aerated biological filter. *Bioresource technology*, 254, 67–74.
- Zhang, Y., Ding, H., Zheng, X., Cai, Z., Misselbrook, T., Carswell, A., & Zhang, J. (2018). Soil N transformation mechanisms can effectively conserve N in soil under saturated conditions compared to unsaturated conditions in subtropical China. *Biology and Fertility of Soils*, 54(4), 495–507.
- Zhang, Y., Wang, X., Hu, M., & Li, P. (2015). Effect of hydraulic retention time (HRT) on the biodegradation of trichloroethylene wastewater and anaerobic bacterial community in the UASB reactor. *Applied microbiology and biotechnology*, 99(4), 1977–1987.
- Zhao, Z., Luo, J., Jin, B., Zhang, J., Li, B., Ma, B., & Shan, B. (2018). Analysis of Bacterial Communities in Partial Nitrification and Conventional Nitrification Systems for Nitrogen Removal. *Scientific reports*, 8.
- Zhu, L., Dong, D., Hua, X., Xu, Y., Guo, Z., & Liang, D. (2017). Ammonia nitrogen removal and recovery from acetylene purification wastewater by air stripping. *Water Science and Technology*, 75(11), 2538-2545.